

Anti-Collagen I Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R013724

Product Information


Application	IHC-P, ELISA, IF (Tissue-P)
Reactivity	Human
Dilution	IHC-P 1:200; IF 1:100
Host	Rabbit
Clonality	Monoclonal
Clone No.	29E23D18
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human Collagen I
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at 20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Collagen I Rabbit mAb [29E23D18] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Synonyms	OI1, OI2, OI3, OI4, EDSC.
Primary Accession	P02452
Gene ID	1277
Background	<p>This gene encodes the pro-$\alpha 1$ chains of type I collagen whose triple helix comprises two $\alpha 1$ chains and one $\alpha 2$ chain. Type I is a fibril-forming collagen found in most connective tissues and is abundant in bone, cornea, dermis and tendon. Mutations in this gene are associated with osteogenesis imperfecta types I-IV, Ehlers-Danlos syndrome type VIIA, Ehlers-Danlos syndrome Classical type, Caffey Disease and idiopathic osteoporosis. Reciprocal translocations between chromosomes 17 and 22, where this gene and the gene for platelet-derived growth factor beta are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans, resulting from unregulated expression of the growth factor. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene. [provided by R. Dalgleish, Feb 2008]</p>




Immunohistochemistry - Anti-Collagen I Rabbit mAb [29E23D18] Sample: Paraformaldehyde-fixed, paraffin embedded human cervical cancer tissue Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins. Primary antibody: R013724 at 1:200 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen. Counter stained with hematoxylin. Positive/negative staining were presented. Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Collagen I Rabbit mAb [29E23D18] Sample: Paraformaldehyde-fixed, paraffin embedded human cervical cancer tissue Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins. Primary antibody: R013724 at 1:200 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen. Counter stained with hematoxylin. Positive/negative staining were presented. Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Collagen I Rabbit mAb [29E23D18] Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins. Primary antibody: R013724 at 1:200 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen. Counter stained with hematoxylin. Positive/negative staining were presented. Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-Collagen I Rabbit mAb [29E23D18] Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins. Primary antibody: R013724 at 1:200 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution DAB was used as the chromogen. Counter stained with hematoxylin. Positive/negative staining were presented. Only the secondary antibody was used as the negative control.

